




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : C07C 237/22, 259/06, C07D 213/56 A61K 31/16, 31/44	A1	(11) International Publication Number: WO 93/24449 (43) International Publication Date: 9 December 1993 (09.12.93)
(21) International Application Number: PCT/GB93/01186 (22) International Filing Date: 3 June 1993 (03.06.93) (30) Priority data: 9211706.8 3 June 1992 (03.06.92) GB (71) Applicant (for all designated States except US): CELLTECH LIMITED [GB/GB]; 216 Bath Road, Slough, Berkshire SL1 4EN (GB). (72) Inventors; and (75) Inventors/Applicants (for US only) : PORTER, John, Robert [GB/GB]; 6 Bader Gardens, Slough, Berkshire SL1 9EN (GB). MORPHY, John, Richard [GB/GB]; 27 Fomersby Crescent, Maidenhead, Berkshire SL6 1YY (GB). MILICAN, Thomas, Andrew [GB/GB]; 3 Harcourt Reach, Dorney Reach, Maidenhead, Berkshire SL6 0DY (GB). BEELEY, Nigel, Robert, Arnold [GB/GB]; 16 Cheshire Road, Thame, Oxfordshire OX9 3LQ (GB).		(74) Agent: HALLYBONE, Huw, George; Carpmaels & Ransford, 43 Bloomsbury Square, London WC1A 2RA (GB). (81) Designated States: AU, CA, JP, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i>
(54) Title: PEPTIDYL DERIVATIVES AND THEIR USE AS METALLOPROTEINASES INHIBITORS <div style="text-align: center;">  <p>(1)</p> </div> (57) Abstract <p>Compounds of formula (1) are described wherein R represents a -CONHOH, carboxyl (-CO₂H), esterified carboxyl or -P(O)(X¹R⁸)X²R⁹ group, where X¹ and X², which may be the same or different is each an oxygen or a sulphur atom, and R⁸ and R⁹, which may be the same or different each represents a hydrogen atom or an optionally substituted alkyl, aryl or aralkyl group; R¹ represents a hydrogen atom or an optionally substituted alkyl, alkenyl, aryl, aralkyl, heteroaralkyl or heteroarylthioalkyl group; R² represents an optionally substituted aryloxy, arylthio, aryloxyalkyl, arylthioalkyl, heteroaryl, heteroaralkyl, heteroaryloxy, heteroarylthio, heteroaryloxyalkyl or heteroarylthioalkyl group; R³ represents a hydrogen atom or an alkyl group; R⁴ represents a hydrogen atom or an alkyl group; R⁵ represents an optionally substituted alkyl or alkenyl group optionally interrupted by one or more -O- or -S- atoms or -N(R⁷) groups, [where R⁷ is a hydrogen atom or a C₁₋₆ alkyl group] or a group -[Alk]_nR⁶ where Alk is an alkyl or alkenyl group optionally interrupted by one or more -O- or -S- atoms or -N(R⁷) groups, n is zero or an integer 1 and R⁶ is an optionally substituted cycloalkyl or cycloalkenyl group; X represents an amino (-NH₂), or substituted amino, hydroxyl or substituted hydroxyl group; and the salts, solvates and hydrates thereof. The compounds are metalloproteinase inhibitors and in particular are selective inhibitors of the action of gelatinase, and may be of use in the treatment of cancer to control the development of tumour metastases.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	MR	Mauritania
AU	Australia	GA	Gabon	MW	Malawi
BB	Barbados	GB	United Kingdom	NL	Netherlands
BE	Belgium	GN	Guinea	NO	Norway
BF	Burkina Faso	GR	Greece	NZ	New Zealand
BG	Bulgaria	HU	Hungary	PL	Poland
BJ	Benin	IE	Ireland	PT	Portugal
BR	Brazil	IT	Italy	RO	Romania
CA	Canada	JP	Japan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SK	Slovak Republic
CI	Côte d'Ivoire	LJ	Liechtenstein	SN	Senegal
CM	Cameroon	LK	Sri Lanka	SU	Soviet Union
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	MC	Monaco	TG	Togo
DE	Germany	MG	Madagascar	UA	Ukraine
DK	Denmark	ML	Mali	US	United States of America
ES	Spain	MN	Mongolia	VN	Viet Nam
FI	Finland				

PEPTIDYL DERIVATIVES AND THEIR USE AS
METALLOPROTEINASES INHIBITORS

Field of the Invention

This invention relates to a novel class of peptidyl derivatives, to processes for their preparation and to their use in medicine.

Background to the Invention

In normal tissues, cellular connective tissue synthesis is offset by extracellular matrix degradation, the two opposing effects existing in dynamic equilibrium. Degradation of the matrix is brought about by the action of proteinases released from resident connective tissue cells and invading inflammatory cells, and is due, in part, to the activity of at least three groups of metalloproteinases. These are the collagenases, the gelatinases (or type-IV collagenases) and the stromelysins. Normally these catabolic enzymes are tightly regulated at the level of their synthesis and secretion and also at the level of their extracellular activity, the latter through the action of specific inhibitors, such as α_2 -macroglobulins and TIMP (tissue inhibitor of metalloproteinase), which form inactive complexes with metalloproteinases.

The accelerated, uncontrolled breakdown of connective tissues by metalloproteinase catalysed resorption of the extracellular matrix is a feature of many pathological conditions, such as rheumatoid arthritis, corneal, epidermal or gastric ulceration; tumour metastasis or invasion; periodontal disease and bone disease. It can be expected that the pathogenesis of such diseases is likely to be modified in a beneficial manner by the administration of metalloproteinase inhibitors and numerous compounds have been suggested for this purpose [for a general review see Wahl, R.C. *et al* Ann. Rep. Med. Chem. 25, 175-184, Academic Press Inc., San Diego (1990)].


Certain hydroxamic acid peptidyl derivatives [see for example European Patent Specifications Nos. 214639, 231081, 236872 and 274453 and International Patent Specifications Nos. WO90/05716 and WO90/05719], have been described as collagenase and/or stromelysin inhibitors.

Summary of the Invention

We have now found a new class of peptidyl derivatives, members of which are metalloproteinase inhibitors and which, in particular, advantageously possess a potent and selective inhibitory action against gelatinase.

There is now much evidence that metalloproteinases are important in tumour invasion and metastasis. Tumour cell gelatinase, in particular, has been associated with the potential of tumour cells to invade and metastasise. Tumour invasion and metastasis is the major cause of treatment failure for cancer patients, and the use of a selective gelatinase inhibitor such as a compound of the present invention which is capable of inhibiting tumour cell invasion can be expected to improve the treatment of this disease.

Thus according to one aspect of the invention we provide a compound of formula (1)



wherein R represents a -CONHOH, carboxyl (-CO₂H), esterified carboxyl or -P(O)(X¹R⁸)X²R⁹ group, where X¹ and X², which may be the same or different is each an oxygen or a sulphur atom, and R⁸ and R⁹, which may be the same or different each represents a hydrogen atom or an optionally substituted alkyl, aryl, or aralkyl group.

R¹ represents a hydrogen atom or an optionally substituted alkyl, alkenyl, aryl, aralkyl, heteroaralkyl or heteroarylthioalkyl group;

R² represents an optionally substituted aryloxy, arylthio, aryloxyalkyl, arylthioalkyl, heteroaryl, heteroaralkyl, heteroaryloxy, heteroarylthio, heteroaryloxyalkyl or heteroarylthioalkyl group;

R³ represents a hydrogen atom or an alkyl group;


R⁴ represents a hydrogen atom or an alkyl group;

R⁵ represents an optionally substituted alkyl or alkenyl group optionally interrupted by one or more -O- or -S- atoms or -N(R⁷)- groups, [where R⁷ is a hydrogen atom or a C₁₋₆alkyl group] or a group -[Alk]_nR⁶ where Alk is an alkyl or alkenyl group optionally interrupted by one or more -O- or -S- atoms or -N(R⁷)- groups, n is zero or an integer 1, and R⁶ is an optionally substituted cycloalkyl or cycloalkenyl group;

X represents an amino (-NH₂), or substituted amino, hydroxyl or substituted hydroxyl group;

and the salts, solvates and hydrates thereof.

It will be appreciated that the compounds according to the invention can contain one or more asymmetrically substituted carbon atoms, for example those marked with an asterisk in formula (1). The presence of one or more of these asymmetric centres in a compound of formula (1) can give rise to stereoisomers, and in each case the invention is to be understood to extend to all such stereoisomers, including enantiomers and diastereoisomers, and mixtures, including racemic mixtures, thereof.

In the formulae herein, the ~line is used at a potential asymmetric centre to represent the possibility of R- and S- configurations, the  line and the ----- line to represent a unique configuration at an asymmetric centre.

In the compounds according to the invention, when the group R represents an esterified carboxyl group, it may be for example a group of formula -CO₂R¹⁵ where R¹⁵ is a straight or branched, optionally substituted C₁₋₈alkyl group such as a methyl, ethyl, n-propyl, i-propyl, n-butyl, i-butyl, s-butyl or t-butyl group; a C₆₋₁₂arylC₁₋₈alkyl group such as an optionally substituted benzyl, phenylethyl, phenylpropyl, 1-naphthylmethyl or 2-naphthylmethyl group; a C₆₋₁₂aryl group such as an optionally substituted

phenyl, 1-naphthyl or 2-naphthyl group; a C₆₋₁₂aryloxyC₁₋₈alkyl group such as an optionally substituted phenyloxymethyl, phenyloxyethyl, 1-naphthyloxymethyl or 2-naphthyloxymethyl group; an optionally substituted C₁₋₈alkanoyloxyC₁₋₈alkyl group, such as a pivaloyloxymethyl, propionyloxyethyl or propionyloxypropyl group; or a C₆₋₁₂aroxyloxyC₁₋₈alkyl group such as an optionally substituted benzoyloxyethyl or benzoyloxypropyl group. Optional substituents present on the groups R¹⁵ include for example one or more halogen atoms such as fluorine, chlorine, bromine or iodine atoms, or C₁₋₄alkyl, e.g. methyl or ethyl, or C₁₋₄alkoxy, e.g. methoxy or ethoxy, groups.

In general, when the group R represents an esterified carboxyl group, it may be a metabolically labile ester of a carboxylic acid.

When the group R¹ in compounds of formula (1) represents an optionally substituted alkyl or alkenyl group, it may be, for example, a straight or branched C₁₋₆ alkyl or C₂₋₆alkenyl group, such as a methyl, ethyl, n-propyl, i-propyl, n-butyl, i-butyl, s-butyl, t-butyl, n-pentyl, i-pentyl, n-hexyl, ethenyl, 1-propenyl, 1-butenyl or 2-butenyl group optionally substituted by one or more C₁₋₆alkoxy, e.g. methoxy, ethoxy, propoxy, C₁₋₆alkylthio, e.g. methylthio, ethylthio, propylthio, C₆₋₁₂arylC₁₋₆alkoxy, e.g. phenylC₁₋₆alkoxy such as benzyloxy, aralkylthio, e.g. phenylC₁₋₆alkylthio such as benzylthio, amino (-NH₂), substituted amino, [such as -NHR¹⁶, where R¹⁶ is a C₁₋₆ alkyl e.g. methyl or ethyl, C₆₋₁₂arylC₁₋₆alkyl, e.g. phenylC₁₋₆alkyl, such as benzyl, C₆₋₁₂aryl, e.g. phenyl, C₃₋₈cycloalkyl, e.g. cyclohexyl, or C₃₋₈cycloalkylC₁₋₆alkyl, e.g. cyclohexylmethyl group], carboxyl (-CO₂H) or -CO₂R¹⁵ [where R¹⁵ is as defined above] groups.

Aryl groups represented by R¹ in compounds of formula (I) include C₆₋₁₂ aryl groups such as phenyl or 1- or 2-naphthyl groups.

Aralkyl groups represented by R¹ include C₆₋₁₂arylC₁₋₆alkyl groups such as phenylC₁₋₆alkyl, or 1- or 2-naphthylC₁₋₆alkyl, for example benzyl, phenylethyl, phenylpropyl, phenylbutyl, phenylpentyl, 1- or 2-naphthylmethyl, naphthylethyl, naphthylpropyl, naphthylbutyl or naphthylpentyl groups.

When the group R¹ in compounds of formula (1) is a heteroaralkyl group, it may be for example a C₃₋₆heteroarylC₁₋₆alkyl group, such as an optionally substituted pyrrolylmethyl, furanylmethyl, thienylmethyl, imidazolylmethyl, oxazolylmethyl, thiazolylmethyl, pyrazolylmethyl, pyridinylmethyl, or pyrimidinylmethyl group.

Heteroarylthioalkyl groups represented by R¹ include C₃₋₆heteroarylthioC₁₋₆alkyl groups such as optionally substituted pyrrolylthiomethyl, furanylthiomethyl, oxazolylthiomethyl, thiazolylthiomethyl, pyrazolylthiomethyl, pyridinylthiomethyl or pyrimidinylthiomethyl groups.

Optional substituents which may be present on heteroaralkyl or heteroarylthioalkyl groups represented by R¹ include those R¹⁰ substituents discussed below.

When the group R² in the compounds according to the invention is an aryloxy, arylthio, aryloxyalkyl or arylthioalkyl group, the aryl portion of the group may be a C₆₋₁₂aryl group such as an optionally substituted phenyl, 1-naphthyl or 2-naphthyl group. The alkyl portion of such groups may be a straight or branched C₁₋₆alkyl group such as a methyl, ethyl, n-propyl, i-propyl, n-butyl, i-butyl, s-butyl, t-butyl, n-pentyl, i-pentyl or n-hexyl group.

When R² is a heteroaryl, heteroaralkyl, heteroaryloxy, heteroarylthio, heteroaryloxyalkyl or heteroarylthioalkyl group, the heteroaryl portion of each group may be a C₃₋₆heteroaryl group containing one, two or three heteroatoms, selected from -O- or -S-, or -N(R⁷)- groups. Particular examples include optionally substituted pyrrolyl, furanyl, thienyl, imadazolyl, oxazolyl, thiazolyl, pyrazolyl, pyridinyl or pyrimidinyl groups. It will be appreciated that such groups may be connected to the remainder of the compound of formula (1) through any ring carbon atom or where appropriate through a heteroatom or group -N(R⁷)-. The alkyl portion of any heteroaryl containing R² group may be a straight or branched C₁₋₆ group, e.g. a C₃₋₆alkyl group, such as a methyl, ethyl, n-propyl, i-propyl, n-butyl, i-butyl, s-butyl, t-butyl, n-pentyl, i-pentyl or n-hexyl group.

The aryl, aralkyl, aryloxy, arylthio, aryloxyalkyl, arylthioalkyl, heteroaryl, heteroaralkyl, heteroaryloxy, heteroarylthio, heteroaryloxyalkyl or heteroarylthioalkyl groups represented by R^1 and/or R^2 in compounds of formula (1) may each optionally be substituted in the cyclic part of the group by one, two or more substituents [R^{10}] selected from halogen atoms, e.g. fluorine, chlorine, bromine or iodine atoms, or C_{1-6} alkyl, e.g. methyl or ethyl, C_{1-6} alkoxy e.g. methoxy or ethoxy, C_{2-6} alkylenedioxy, e.g. ethylenedioxy, halo C_{1-6} alkyl, e.g. tri-fluoromethyl, C_{1-6} alkylamino, e.g. methylamino or ethylamino, C_{1-6} dialkylamino, e.g. dimethylamino or diethylamino, amino ($-NH_2$), nitro, cyano, hydroxyl ($-OH$), carboxyl ($-CO_2H$), $-CO_2R^{15}$, where R^{15} is as defined above, C_{1-6} alkylcarbonyl, e.g. acetyl, sulphonyl ($-SO_3H$), C_{1-6} alkylsulphonyl, e.g. methylsulphonyl, aminosulphonyl ($-SO_2NH_2$), C_{1-6} alkylaminosulphonyl, e.g. methylaminosulphonyl or ethylaminosulphonyl, C_{1-6} dialkylaminosulphonyl e.g. dimethylaminosulphonyl or diethylaminosulphonyl, carboxamido ($-CONH_2$), C_{1-6} alkylaminocarbonyl, e.g. methylaminocarbonyl or ethylaminocarbonyl, C_{1-6} dialkylaminocarbonyl, e.g. dimethylaminocarbonyl or diethylaminocarbonyl, sulphonylamino ($-NHSO_2H$), C_{1-6} alkylsulphonylamino, e.g. methylsulphonylamino or ethylsulphonylamino, C_{1-6} dialkylsulphonylamino, e.g. dimethylsulphonylamino or diethylsulphonylamino, C_{1-6} alkanoylamino, e.g. acetylamino or C_{1-6} alkanoylamino C_{1-6} alkyl e.g. acetylaminomethyl groups. It will be appreciated that where two or more R^{10} substituents are present, these need not necessarily be the same atoms and/or groups. The R^{10} substituents may be present at any ring carbon atom away from that attached to the rest of the molecule of formula (1). Thus, for example, in phenyl groups any substituents may be present at the 2-, 3- or 4- 5- or 6-positions relative to the ring carbon atom attached to the remainder of the molecule.

When the groups R^3 and R^4 in compounds of formula (1) are alkyl groups, they may be for example C_{1-6} alkyl groups such as methyl or ethyl groups.

The group R^5 in compounds of formula (1) may be an optionally substituted straight or branched C_{1-6} alkyl, e.g. methyl, ethyl, n-propyl, i-propyl, n-butyl, i-butyl, t-butyl, n-pentyl or n-hexyl or C_{2-6} alkenyl e.g. ethenyl or 1-propenyl group optionally interrupted by one or more $-O-$ or $-S-$ atoms or $-N(R^7)-$

groups where R^7 is a hydrogen atom or a C_{1-6} alkyl group such as a methyl group.

Optional substituents which may be present on alkyl or alkenyl groups R^5 include C_{6-12} aryl C_{1-6} alkyl groups such as optionally substituted phenyl C_{1-6} alkyl e.g. benzyl groups, C_{6-12} aryl C_{1-6} alkoxy groups such as optionally substituted phenyl C_{1-6} alkoxy e.g. benzyloxy groups, C_{6-12} aryl e.g. optionally substituted phenyl groups, C_{3-8} heteroaryl e.g. optionally substituted indole, imidazole or quinoline groups, C_{6-12} aryl C_{1-6} alkoxy C_{6-12} aryl, e.g. benzyloxyphenyl groups, -OH, -SH, C_{1-6} alkylthio e.g. methylthio or ethylthio, carboxy (-CO₂H), amino (-NH₂), carboxamido (-CONH₂) or guanido -NHC(NH₂)=NH, groups. The optional substituents present on these groups may be R^{10} substituents as discussed above.

When the group Alk is present in compounds of formula (1) it may be a straight or branched C_{1-6} alkyl, e.g. methyl, ethyl, n-propyl i-propyl, n-butyl, i-butyl, t-butyl, n-pentyl or n-hexyl or C_{2-6} alkenyl e.g. ethenyl or 1-propenyl group optionally interrupted by one or more -O- or -S- atoms or -N(R^7)- groups where R^7 is a hydrogen atom or a C_{1-6} alkyl group such as a methyl group.

The group R^6 in compounds of formula (1) may represent a C_{3-8} cycloalkyl, e.g. cyclopentyl or cyclohexyl, or C_{3-8} cycloalkenyl e.g. cyclopentenyl or cyclohexenyl, group optionally substituted by one, two or more C_{1-6} alkyl, e.g. methyl or ethyl, C_{1-6} alkoxy, e.g. methoxy or ethoxy, C_{1-6} alkylthio, e.g. methylthio, or hydroxyl groups.

When X in the compounds of formula (1) represents a substituted amino group it may be for example a group of formula -NR¹¹R¹², where R^{11} and R^{12} , which may be the same or different, is each a hydrogen atom (with the proviso that when one of R^{11} or R^{12} is a hydrogen atom, the other is not) or an optionally substituted straight or branched alkyl group, optionally interrupted by one or more -O- or -S- atoms or -N(R^7)- or aminocarbonyloxy [-NHC(O)O-] groups or R^{11} and R^{12} , together with the nitrogen atom to which they are attached, may form an optionally substituted C_{3-6} cyclic amino group optionally possessing one or more other heteroatoms selected

from -O-, -S- or -N(R)⁷- groups.

When R¹¹ and/or R¹² is an alkyl group it may be for example a C₁₋₆alkyl group such as a methyl, ethyl, n-propyl, i-propyl, n-butyl, i-butyl, s-butyl, or t-butyl group, optionally interrupted by one or more -O- or -S- atoms, or -N(R⁷)- or aminocarbonyloxy groups and may be for example a methoxymethyl, ethoxymethyl, methoxyethyl, ethoxyethyl or ethylaminocarbonyloxymethyl group. The optional substituents which may be present on such groups include hydroxyl (-OH), carboxyl (-CO₂H), esterified carboxyl (-CO₂R¹⁵), carboxamido (-CONH₂), substituted carboxamido, e.g. a group -CONR¹¹R¹² where NR¹¹R¹² is as defined herein, amino (-NH₂), substituted amino, for example a group of formula -NR¹¹R¹², aminosulphonylamino, for example -N(R⁷)SO₂NH₂ or -N(R⁷)SO₂NR¹¹R¹², or aryl, e.g. C₆₋₁₂ aryl such as phenyl, optionally substituted by one, two or more R¹⁰ substituents selected from those listed above in relation to the group R².

Particular examples of cyclic amino groups represented by -NR¹¹R¹² include morpholinyl, imidazolyl, piperazinyl, pyrrolyl, oxazolyl, thiazolyl, pyrazolyl, pyrrolidinyl, pyridinyl and pyrimidinyl groups.

The groups R⁸ or R⁹ in compounds of formula (1) may be hydrogen atom or an optionally substituted straight or branched C₁₋₆alkyl, e.g. methyl, ethyl, n-propyl, i-propyl, n-butyl or i-butyl, C₆₋₁₂aryl e.g. phenyl, or C₆₋₁₂arylC₁₋₆alkyl, e.g. benzyl, phenylethyl or phenylpropyl group. Optional substituents present on alkyl groups of this type include one or more C₁₋₆alkoxy, e.g. methoxy, ethoxy, or C₁₋₆alkylthio e.g. methylthio or ethylthio groups or an optionally substituted C₆₋₁₂aryloxy, e.g. phenyloxy, C₆₋₁₂arylthio e.g. phenylthio, C₆₋₁₂arylC₁₋₆alkoxy e.g. benzyloxy or C₆₋₁₂arylC₁₋₆alkylthio e.g. benzylthio. Optional substituents present on the groups R⁸ or R⁹ when it is an aryl or aralkyl group or an alkyl group substituted by an aryloxy or arylthio group include R¹⁰ groups present on the cyclic part of R⁸ or R⁹ as defined above.

Salts of compounds of formula (1) include pharmaceutically acceptable salts, for example acid addition salts derived from inorganic or organic

acids, such as hydrochlorides, hydrobromides, hydroiodides, p-toluene sulphonates, phosphates, sulphates, acetates, trifluoroacetates, propionates, citrates, maleates, fumarates, malonates, succinates, lactates, oxalates, tartarates and benzoates.

Salts may also be formed with bases. Such salts include salts derived from inorganic or organic bases, for example alkali metal salts such as sodium or potassium salts, alkaline earth metal salts such as magnesium or calcium salts, and organic amine salts such as morpholine, piperidine, dimethylamine or diethylamine salts.

When the group R in compounds of the invention is an esterified carboxyl group, it may be a metabolically labile ester of formula $-\text{CO}_2\text{R}^{15}$ where R^{15} may be an ethyl, benzyl, phenylethyl, phenylpropyl, 1- or 2-naphthyl, 2,4-dimethylphenyl, 4-t-butylphenyl, 2,2,2-trifluoroethyl, 1-(benzyloxy)benzyl, 1-(benzyloxy)ethyl, 2-methyl-1-propionyloxypropyl, 2,4,6-trimethylbenzoyloxymethyl or pivaloyloxymethyl group.

When the group R in compounds of formula (1) is a $-\text{P}(\text{O})(\text{X}^1\text{R}^8)\text{X}^2\text{R}^9$ group it may in particular be a $-\text{P}(\text{O})(\text{OH})\text{OR}^9$, $-\text{P}(\text{O})(\text{SH})\text{OR}^9$ or $-\text{P}(\text{O})(\text{OH})\text{SR}^9$ group. Examples of such groups include $-\text{P}(\text{O})(\text{OH})\text{OH}$, $-\text{P}(\text{O})(\text{OH})\text{SH}$, $-\text{P}(\text{O})(\text{SH})\text{OH}$, $-\text{P}(\text{O})(\text{OH})\text{OCH}_3$, $-\text{P}(\text{O})(\text{OH})\text{SCH}_3$, $-\text{P}(\text{O})(\text{OH})\text{OCH}_2\text{CH}_3$, $-\text{P}(\text{O})(\text{OH})\text{OPh}$, $-\text{P}(\text{O})(\text{OH})\text{SPh}$, $-\text{P}(\text{O})(\text{OH})\text{OCH}_2\text{Ph}$ or $-\text{P}(\text{O})(\text{OH})\text{SCH}_2\text{Ph}$, where Ph is a phenyl group optionally substituted by one or more substituents R^{10} .

In the compounds of formula (1) the group R^1 may in particular be a C_{1-6} alkyl group such as a methyl group, an aralkyl group such as benzyl group, an arylthioalkyl group such as a phenylthiomethyl group or a heteroarylthioalkyl group such as thienylthiomethyl, pyridinylthiomethyl or pyrimidinylthiomethyl group or is especially a hydrogen atom.

The group R^2 may be in particular an optionally substituted phenoxy, phenylthio, phenoxyalkyl, e.g. phenoxyethyl, phenoxyethyl, phenoxypropyl, phenoxybutyl, phenylthioalkyl, e.g. phenylthiomethyl, phenylthioethyl, phenylthiopropyl, or phenylthiobutyl group or a group Het-

Het-oxy, Het-thio, Het-propyl, Het-butyl, Het-pentyl, Het-oxymethyl, Het-oxyethyl, Het-oxypropyl, Het-oxybutyl, Het-thiomethyl, Het-thioethyl, Het-thiopropyl or Het-thiobutyl, wherein Het is an optionally substituted pyrrolyl, furanyl, thienyl, imadazolyl, oxazolyl, thiazolyl, pyrazolyl, pyrimidinyl or, especially, pyridinyl group connected to the remainder of the group through a ring carbon atom or a heteroatom or -N(R⁷) group. Optional substituents present on R² groups of the above types include one, two or more R¹⁰ substituents as defined above. Where the group is an optionally substituted phenyl group, it is preferably a 4-substituted phenyl group, for example a 4-halophenyl, e.g. 4-chlorophenyl, 4-alkoxyphenyl, e.g. 4-methoxyphenyl or 4-alkylphenyl, e.g. 4-methylphenyl group.

Particularly useful R² groups are aryloxyalkyl, heteroaralkyl or heteroraryloxyalkyl groups. Preferred groups of these types are optionally substituted phenoxyalkyl, e.g. phenoxyethyl, phenoxypropyl, or phenoxybutyl groups, Het-oxy C₁₋₆ alkyl such as pyridinyl-oxyethyl, pyridinyl-oxypropyl or pyridinyl-oxybutyl groups, or C₃₋₆ heteroaryl C₁₋₆ alkyl, e.g. C₃₋₆ heteroaryl C₃₋₆ alkyl such as C₃₋₆ heteroarylpropyl groups.

The groups R³ and R⁴ in compounds of formula (1) may each in particular be a methyl group, or, especially, a hydrogen atom.

The group R⁵ in compounds of formula (1) may be in particular a group -AlkR⁶, where R⁶ is an optionally substituted cycloalkyl or cycloalkenyl group.

Thus, the group R⁵ in compounds of formula (1) may be an optionally substituted C₃₋₈cycloalkylC₁₋₆alkyl [e.g. cyclopentylC₁₋₆alkyl such as cyclopentylmethyl or cyclopentylethyl, or cyclohexylC₁₋₆alkyl such as cyclohexylmethyl or cyclohexylethyl], C₃₋₈cycloalkenylC₁₋₆alkyl [e.g. cyclopentenylC₁₋₆alkyl such as cyclopentenylmethyl or cyclohexenylC₁₋₆alkyl such as cyclohexenylmethyl], cycloalkylC₁₋₃alkoxyC₁₋₃alkyl [e.g. cyclopentylmethoxymethyl, cyclohexylmethoxymethyl] C₃₋₈cycloalkenylC₁₋₃alkoxyC₁₋₃alkyl [e.g. cyclopentenylmethoxymethyl or cyclohexenylmethoxymethyl] C₃₋₈cycloalkylC₁₋₃alkylthioC₁₋₃alkyl [e.g. cyclopentylmethylthiomethyl or cyclohexylmethylthiomethyl] or C₃₋

$_{8\text{cycloalkenylC}_{1-3}\text{alkylthioC}_{1-3}\text{alkyl}}$ [e.g. cyclopentenylmethylthiomethyl or cyclohexenylmethylthiomethyl], $_{\text{C}_{3-8}\text{cycloalkylC}_{1-3}\text{alkylaminoC}_{1-3}\text{alkyl}}$ [e.g. cyclopentylmethylaminomethyl, or cyclohexylmethylaminomethyl] or $_{\text{C}_{3-8}\text{cycloalkenylC}_{1-3}\text{alkylaminoC}_{1-3}\text{alkyl}}$ [e.g. cyclopentenylmethylaminomethyl or cyclohexenylmethylaminomethyl] group.

In another preference, the group R^5 in compounds of formula (1) may in particular be a C_{1-6} alkyl group, e.g. an i-propyl, i-butyl or t-butyl group, or an optionally substituted benzyl, benzyloxybenzyl or indolylmethyl group.

The group X in compounds of formula (1) may be in particular an amino ($-\text{NH}_2$) or $-\text{NR}^{11}\text{R}^{12}$ group. Particular $-\text{NR}^{11}\text{R}^{12}$ groups are $-\text{NHR}^{12}$ groups. Groups of this type include those where R^{12} is a C_{1-6} alkyl group, for example a methyl, ethyl, or n-propyl group, optionally interrupted by one or more -O- or -S- atoms or $-\text{N}(\text{R}^7)$ [e.g. $-\text{NH}-$ or $-\text{N}(\text{CH}_3)-$] or aminocarbonyloxy groups and optionally substituted by a hydroxyl, carboxyl, carboxyalkyl, e.g. carboxymethyl, carboxamido, amino, $-\text{NR}^{11}\text{R}^{12}$, [for example di- C_{1-6} alkylamino such as dimethylamino, C_{1-6} alkylamino such as methylamino, or C_{3-6} cyclic amino such as morpholinyl, pyrrolidinyl or pyridinyl] aminosulphonylamino [for example $-\text{N}(\text{R}^7)\text{SO}_2\text{NH}_2$ such as $-\text{NHSO}_2\text{NH}_2$ or $-\text{N}(\text{CH}_3)_3\text{SO}_2\text{NH}_2$ or $-\text{N}(\text{R}^7)\text{SO}_2\text{NR}^{11}\text{R}^{12}$ such as $-\text{NHSO}_2\text{R}^{11}\text{R}^{12}$ or $-\text{N}(\text{CH}_3)\text{SO}_2\text{NR}^{11}\text{R}^{12}$, especially where R^{11} and R^{12} is each an optionally substituted C_{1-6} alkyl group or R^{11} and R^{12} together with the nitrogen atom to which they are attached is an optionally substituted C_{3-6} cyclic amino group] or phenyl optionally substituted by one, two or more R^{10} substituents.

A particularly useful group of compounds according to the invention is that of formula (1) wherein R^5 is an AlkR^6 group, where Alk is a C_{1-6} alkyl and R^6 is a cycloalkyl or cycloalkenyl group.


In another preference, compounds of formula (1) wherein R^5 is a propyl, e.g. i-propyl or butyl, e.g. i-butyl or especially t-butyl group are particularly useful.

In general, in compounds of formula (1) the groups R^1 , R^3 and R^4 is each preferably a hydrogen atom.

In a further preference, the group R in compounds according to the invention is a -CONHOH or a -CO₂H group or a metabolically labile ester thereof, or a group P(O)(OH)OR⁷. In a particular preference, however, R is a -CO₂H or -P(O)(OH)₂ group or especially a -CONHOH group.

X in the compounds of formula (1) is preferably an amino or substituted amino group.

An especially useful group of compounds according to the invention has the formula (1a):



wherein R, R², R⁵ and X are as defined for formula (1); and the salts, solvates and hydrates thereof.

It will be appreciated that the particular definitions and preferences recited above with respect to the groups R, R², R⁵ and X also apply to these groups in compounds of formula (1a). In addition, a particularly useful group of compounds of formula (1a) are those wherein R represents a -CONHOH, or -CO₂H or -P(O)(OH)₂ group; R² is as defined for formula (1);

R⁵ represents a group -AlkR⁶, where Alk is a C₁₋₆ alkyl group and R⁶ is a cycloalkyl or cycloalkenyl group or a C₁₋₆ alkyl group;

X is an amino or substituted amino group; and the salts, solvates and hydrates thereof.

Particularly useful compounds of formula (1a) are those wherein R⁵ is a group -AlkR⁶, and R⁶ is an optionally substituted cyclohexyl group. Compounds of this type in which R⁵ is a cyclohexylC₁₋₆alkyl group, particularly a cyclohexylmethyl group, are especially useful.

In another preference, compounds of formula (1a) wherein R⁵ is a propyl, e.g. i-propyl or especially butyl, e.g. t-butyl, are particularly useful.

In the compounds of formula (1a) X may be a -NH₂ or -NR¹¹R¹² group.

In the compounds of formulae (1) and (1a), when the group R⁵ is a cycloalkylC₁₋₆alkyl group then the chiral centre to which this group is attached preferably has a S-configuration.

Particularly useful compounds according to the invention are:

{4-(N-hydroxyamino)-2(R)-[3-(4-pyridinium)propyl]succinyl}-L-β-cyclohexylalanine-N-(2-phenylethyl)amide;

{4-(N-hydroxyamino)-2(R)-[3-(4-pyridinium)propyl]succinyl}-L-β-cyclohexylalanine-N[2-(p-sulphonamidophenyl)ethyl]amide;


{4-(N-hydroxyamino)-2(R)-[3-(4-pyridinium)propyl]succinyl}-L-t-leucine amide;

{4-(N-hydroxyamino)-2(R)-[3-(2-phenoxyethyl)succinyl]-L-β-cyclohexylalanine-N-(2-phenylethyl)amide; and the salts solvates and hydrates thereof.


The compounds according to the invention may be prepared by the following processes. In the description and formulae below the groups R, R¹, R², R³, R⁴, R⁵ and X are as defined above, except where otherwise indicated. It will be appreciated that functional groups, such as amino, hydroxyl or carboxyl groups, present in the various compounds described below, and which it is desired to retain, may need to be in protected form before any reaction is initiated. In such instances, removal of the protecting group may be the final step in a particular reaction. Suitable amino or hydroxyl protecting groups include benzyl, benzyloxycarbonyl or t-butyloxycarbonyl groups. These may be removed from a protected derivative by catalytic hydrogenation using for example hydrogen in the

presence of a metal catalyst, for example palladium on a support such as carbon in a solvent such as an alcohol e.g. methanol, or by treatment with trimethylsilyl iodide or trifluoroacetic acid in an aqueous solvent. Suitable carboxyl protecting groups include benzyl groups, which may be removed from a protected derivative by the methods just discussed, or alkyl groups, such as a t-butyl group which may be removed from a protected derivative by treatment with trifluoroacetic acid in an aqueous solvent. Other suitable protecting groups and methods for their use will be readily apparent. The formation of the protected amino, hydroxyl or carboxyl group may be achieved using standard alkylation or esterification procedures, for example as described below.

Thus according to a further aspect of the invention a compound of formula (1) may be prepared by coupling an acid of formula (2)



or an active derivative thereof, with an amine of formula (3)



followed by removal of any protecting groups.

Active derivatives of acids for formula (2) include for example acid anhydrides, or acid halides, such as acid chlorides.


The coupling reaction may be performed using standard conditions for amination reactions of this type. Thus, for example the reaction may be

achieved in a solvent, for example an inert organic solvent such as an ether, e.g. a cyclic ether such as tetrahydrofuran, an amide e.g. a substituted amide such as dimethylformamide, or a halogenated hydrocarbon such as dichloromethane at a low temperature, e.g. -30°C to ambient temperature, such as -20°C to 0°C , optionally in the presence of a base, e.g. an organic base such as an amine, e.g. triethylamine or a cyclic amine such as N-methylmorpholine. Where an acid of formula (2) is used, the reaction may additionally be performed in the presence of a condensing agent, for example a diimide such as N,N'-dicyclohexylcarbodiimide or 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride, advantageously in the presence of a triazole such as 1-hydroxybenzotriazole. Alternatively, the acid may be reacted with a chloroformate for example ethylchloroformate, prior to reaction with the amine of formula (3).


Free hydroxyl or carboxyl groups in the starting materials of formulae (2) and (3) may need to be protected during the coupling reaction. Suitable protecting groups and methods for their removal may be those mentioned above. Where R in the intermediates of formula (2) is a $-\text{P}(\text{O})(\text{X}^1\text{R}^8)\text{X}^2\text{R}^9$ group, at least one of R^8 or R^9 is other than a hydrogen atom. Conveniently, each of R^8 and R^9 is a optionally substituted alkyl, aryl or aralkyl group. Such groups, when present in compounds of the invention may be cleaved as described below to yield other compounds of the invention wherein R^8 and/or R^9 is each a hydrogen atom.

When a particular stereoisomer of formula (1) is required, the resolution of a mixture of isomer is the same as described in WO 92/09564.

Where desired, however, appropriate homochiral starting materials may be used in the coupling reaction to yield a particular stereo isomer of formula (1). Thus, in a particular process a compound of formula (1a) may be prepared by reaction of a compound of formula (2a)




with an amine of formula (3a)



as described above

Intermediate acids of formula (2) wherein R is a or a group $-P(O)(X^1R^8)X^2R^9$ may be prepared from a corresponding ester of formula (4)



where R^{13} is an alkyl group, for example a methyl or t-butyl group, by hydrolysis using for example trifluoroacetic acid, or, when R^{13} is an aralkyl group, such as a benzyl group, by hydrogenolysis, for example by reaction with hydrogen in the presence of a metal catalyst, e.g. palladium, on a support such as carbon in a solvent such as an alcohol, e.g. methanol optionally at an elevated pressure and temperature.

Intermediate esters of formula (4) where R is a $-P(O)(X^1R^8)X^2R^9$ group may be prepared by reaction of an acrylate $R^1CHC(R^2)COR^{13}$ with a phosphite $-P(OR^{14})(X^1R^8)X^2R^9$ [where R^{14} is a leaving group, for example a silyl group such as a trialkylsilyl group e.g. a trimethylsilyl group] at an elevated temperature.

Acrylates of formula $R^1CHC(R^2)COOR^{13}$ may be prepared by reaction of a mono-ester $HOOCCH(R^2)COOR^{13}$ with an aldehyde R^1CHO or a polymer thereof e.g. paraformaldehyde or paraldehyde in the presence of a base, for


example an organic base such as piperidine. The reaction may be performed in a solvent, such as pyridine, optionally at an elevated temperature.

Mono-esters of formula $\text{HOOCCH(R}^2\text{)COOR}^{13}$ may be prepared by hydrolysis of the corresponding di-ester $\text{R}^{13}\text{OOCCH(R}^2\text{)COOR}^{13}$ using a base, for example an alkali hydroxide, in an inert solvent such as dioxan at a low temperature e.g. around 0°C . The di-esters for use in this reaction may be prepared by alkylation of the corresponding malonates of formula $\text{R}^{13}\text{OOCCH}_2\text{COOR}^{13}$ with a halide R^2Hal [where Hal is a halogen atom such as a chlorine or bromine atom] in the presence of a base, e.g. a hydride such as sodium hydride in a solvent such as tetrahydrofuran at ambient temperature. Malonates of formula $\text{R}^{13}\text{OOCCH}_2\text{COOR}^{13}$ are either known compounds or may be prepared by methods analogous to those used for the preparation of the known compounds.

Intermediate phosphites of formula $-\text{P(OR}^{14}\text{)(X}^1\text{R}^8\text{)X}^2\text{R}^9$ may be prepared by reaction of a phosphite $-\text{P(O)(X}^1\text{R}^8\text{)X}^2\text{R}^9$ with an appropriate amine $(\text{R}^{14})_2\text{NH}$ e.g. a silazane, at an elevated temperature, e.g. the reflux temperature. Phosphites of formula $\text{P(O)(X}^1\text{R}^8\text{)X}^1\text{R}^9$ are either known compounds or may be prepared by methods analogous to those used for the preparation of the known compounds.

Intermediate amines of formula (3) may be prepared by reaction of the corresponding acids $\text{R}^3\text{NHC(R}^4\text{)(R}^5\text{)COOH}$ or active derivatives thereof with an amine XH using the reagents and conditions described above for the preparation of compounds of formula (1) from intermediates for formula (2) and (3). The acids $\text{R}^3\text{NHC(R}^4\text{)(R}^5\text{)COOH}$ and amines XH are either known compounds or may be prepared from known starting materials using analogous processes [for example as described in the preparation of the specific Intermediates in the Examples herein] to those used for the preparation of the known compounds.

The homochiral acids of formula (2a) may be prepared according to another feature of the invention by oxidation of an oxazolidinone of formula (5)



(where Ph is a phenyl group)

using an oxidising agent such as peroxide, e.g. hydrogen peroxide in a solvent such as an ether e.g. a cyclic ether such as tetrahydrofuran, at a low temperature, e.g. around 0°C followed by treatment with a base, such as lithium hydroxide, at an elevated temperature.

The compounds of formula (5) are novel, particularly useful, intermediates for the preparation of stereoisomers of formula (1a) and form a further aspect of the invention.

The compounds of formula (5) may be prepared by reaction of an acyl halide $(R^2)COHal$ (where Hal is a halogen atom such as chlorine, bromine or iodine atom) with a solution of (S)-4-(phenylmethyl)-2-oxazolidinone in the presence of a base such as n-butyl lithium in a solvent such as tetrahydrofuran at a low temperature, e.g. around -78°C, followed by treatment of the resulting oxazolidinone with a reagent RCH_2Hal in the presence of a silazide such as sodium hexamethyldisilazide at a low temperature.

Acyl halides R^2COHal may be prepared by treatment of the corresponding known acids R^2CO_2H with conventional halogenating agents for example thionyl halides such as thionyl chloride under standard reaction conditions.

In an other process, compounds of formula (5) wherein R^2 is a heteroarylpropyl group may be prepared by reaction of an acid $R^2CH_2CO_2H$ (wherein R^2 is as just defined) with a solution of (S)-4-(phenylmethyl)-2-oxazolidinone in the presence of a condensing agent such as dicyclohexylcarbodiimide or N,N-dimethylaminopyridine. The acid itself may be prepared by hydrolysis of the corresponding ester $R^2CH_2CO_2R^{13}$

using well known methods, for example by treatment with lithium hydroxide in methanol at room temperature.

Esters $R^2CH_2CO_2R^{13}$ may be prepared by catalytic reduction of a diene of formula (6)



(wherein Het is a heteroaryl group as defined above for R²) using standard methods, for example by catalytic hydrogenation using for example hydrogen in the presence of a catalyst such as palladium on carbon in a solvent such as methanol.

The diene of formula (6) may be prepared by coupling an aldehyde HetCHO with a phosphonate such as trimethylphosphonocrotonate in the presence of a silazide such as sodium hexamethyldisilazide in tetrahydrofuran at a temperature of about 0°C.

Compounds of formula (1) may also be prepared by interconversion of other compounds of formula (1). Thus, for example, a compound of formula (1) wherein R is a -CONHOH group may be prepared by reaction of a corresponding acid of formula (1) wherein R is a -CO₂H group or an active derivative thereof (for example an acid chloride or an acid anhydride) with hydroxylamine or an O-protected derivative for example O-trimethylsilylhydroxylamine or a salt thereof. The reaction may be performed using the reagents and conditions described above in the preparation of compounds of formula (1) from the starting materials of formulae (2) and (3). If desired the acid starting material may be reacted with a chloroformate, for example ethyl chloroformate, prior to reaction with the hydroxylamine or protected hydroxylamine.

In another interconversion process, compounds of formula (1) wherein R is -CO₂H may be prepared by hydrolysis of the corresponding esterified compounds (for example where R is a -CO₂R¹⁵ group and/or X contains a similar group) using conventional procedures, for example by treatment with

a base, e.g. an alkali metal hydroxide such as lithium hydroxide in a solvent such as an aqueous alcohol, e.g. aqueous methanol, or by treatment with an acid such as a mineral acid, e.g. hydrochloric acid in the presence of a solvent, e.g. dioxan.

Similarly esters of formula (1) for example where R is a CO_2R^{15} group and/or X contains a $-\text{CO}_2\text{R}^{15}$ group may be prepared by reaction of the corresponding acids, where R is a $-\text{CO}_2\text{H}$ group and/or X contains a $-\text{CO}_2\text{H}$ group or an active derivative thereof, with an alcohol R^{15}OH using standard conditions.

The compounds according to the invention are potent and selective inhibitors of gelatinase. The activity and selectivity of the compounds may be determined by the use of appropriate enzyme inhibition test for example as described in Example A hereinafter. In our tests using this approach, compounds according to the invention have been shown to inhibit gelatinase with K_i values in the picomolar-nanomolar range and to have around a 40 fold or greater selectivity for gelatinase over stromelysin, and around a 100-fold or greater selectivity for gelatinase over collagenase.

The ability of compounds of the invention to prevent tumour cell invasion may be demonstrated in a standard mouse model.

Thus, briefly, nude mice may be inoculated with a tumour cell line showing gelatinase - dependent invasion and the ability of compounds according to the invention to reduce subsequent lung tumour colonisation may be evaluated in accordance with standard procedures. In our tests, compounds according to the invention, when administered intravenously at 1mg/kg to mice in the above model have reduced lung tumour colonisation to negligible levels.

The compounds according to the invention can be expected to be of use to prevent tumour cell metastasis and invasion. The compounds may therefore be of use in the treatment of cancer, particularly in conjunction with radiotherapy, chemotherapy or surgery, or in patients presenting with primary tumours, to control the development of tumour metastases. Thus,

according to a further aspect of the invention we provide a compound of formula (1) for use in the treatment of cancer to control the development of tumour metastases. Particular cancers may include breast, melanoma, lung, head, neck or bladder cancers.

For use according to this aspect of the invention, the compounds of formula (1) may be formulated in a conventional manner, optionally with one or more physiologically acceptable carriers, diluents or excipients.

Thus according to a further aspect of the invention we provide a pharmaceutical composition comprising a compound of formula (1) and a pharmaceutically acceptable diluent, carrier or excipient.

In a still further aspect the invention provides a process for the production of a pharmaceutical composition comprising bringing a compound of formula (1) into association with a pharmaceutically acceptable diluent, carrier or excipient.

Compounds for use according to the present invention may be formulated for oral, buccal, parental or rectal administration or in a form suitable for nasal administration or administration by inhalation or insufflation.

For oral administration, the pharmaceutical compositions may take the form of, for example, tablets or capsules prepared by conventional means with pharmaceutically acceptable excipients such as binding agents (e.g. pregelatinised maize starch, polyvinylpyrrolidone or hydroxypropyl methylcellulose); fillers (e.g. lactose, microcrystalline cellulose or calcium hydrogen phosphate); lubricants (e.g. magnesium stearate, talc or silica); disintegrants (e.g. potato starch or sodium glycollate); or wetting agents (e.g. sodium lauryl sulphate). The tablets may be coated by methods well known in the art. Liquid preparations for oral administration may take the form of, for example, solutions, syrups or suspensions, or they may be presented as a dry product for constitution with water or other suitable vehicle before use. Such liquid preparations may be prepared by conventional means with pharmaceutically acceptable additives such as suspending agents, emulsifying agents, non-aqueous vehicles; and preservatives. The

preparations may also contain buffer salts, flavouring, colouring and sweetening agents as appropriate.

Preparations for oral administration may be suitably formulated to give controlled release of the active compound.

For buccal administration the compositions may take the form of tablets or lozenges formulated in conventional manner.

The compounds of formula (1) may be formulated for parental administration by injection e.g. by bolus injection or continuous infusion. Formulations for injection may be presented in unit dosage form. The compositions for injection may take such forms as suspensions, solutions or emulsions in oily or aqueous vehicles, and may contain formulatory agents such as suspending, stabilising and/or dispersing agents. Alternatively, the active ingredient may be in powder form for constitution with a suitable vehicle, e.g. sterile pyrogen-free water, before use.

The compounds of formula (1) may also be formulated in rectal compositions such as suppositories or retention enemas, e.g. containing conventional suppository bases such as cocoa butter or other glycerides.

In addition to the formulations described above the compounds of formula (1) may also be formulated as a depot preparation. Such long acting formulations may be administered by implantation or by intramuscular injection.

For nasal administration or administration by inhalation the compounds for use according to the present invention are conveniently delivered in the form of an aerosol spray presentation for pressurised packs or a nebuliser, with the use of suitable propellant, e.g. dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, carbon dioxide or other suitable gas.

The compositions may, if desired, be presented in a pack or dispenser device which may contain one or more unit dosage forms containing the

active ingredient. The pack or dispenser device may be accompanied by instructions for administration.

The doses of compounds of formula (1) used to control the development of tumour metastases will vary depending on the condition of the patient to be treated but in general may be in the range around 0.5mg to 100mg/kg body weight, particularly from about 1mg to 40mg/kg body weight. Dosage units may be varied according to the route of administration of the compound and condition of the patient in accordance with conventional practice.

The following Examples illustrate the invention.

Example A

The activity and selectivity of the compounds of the invention may be determined as described below.

All enzyme assays to determine K_i values were performed using the peptide substrate Dnp-Pro-Leu-Gly-Trp-Ala-D-Arg-NH₂. [M. Sharon Stock and Rober D. Gray, JBC 264, 4277-81, 1989). The enzymes cleave at the Gly-Leu bond which can be followed fluorimetrically by measuring the increase in Trp fluorescence emission associated with the removal of the quenching dinitrophenol (Dnp) group.

Essentially, enzyme (e.g. gelatinase, stromelysin, collagenase) at 0.08-2nM; a range of inhibitor concentrations (0.1-50 x K_i) and substrate (approx. 20 μ M) are incubated overnight in 0.1M Tris/HCl buffer, pH 7.5, containing 0.1M NaCl, 10mM CaCl₂ and 0.05% Brij 35 at either room temperature or 37°C depending on the enzyme. The reaction is stopped by adjusting the pH to 4 using 0.1M sodium acetate buffer and the fluorescence read at an excitation wavelength of 280nm and emission wavelength of 346nm.

K_i values can be established using the equation for tight-binding inhibition: -

$$V_i = \frac{V_o}{2[E]} \left(\sqrt{(K_{i(app)} + [I])^2 + 2 (K_{i(app)} - [I])[E] + [E]^2} - (K_{i(app)} + [I] - [E]) \right)$$

where V_o is the initial rate of reaction in the absence of inhibitor, V_i is the initial rate in the presence of inhibitor, $[E]$ is the total enzyme concentration and $[I]$ the total inhibitor concentration in the reaction mixture.

For stromelysin and collagenase, K_i (app) was assumed to approximate to the true K_i as $[S] \ll K_m$ for the substrate hydrolysis. For gelatinase the K_i was determined by performing the analyses at several substrate concentrations. A plot of K_i (app) vs. $[S]$ then gave the true K_i as the value of the y-axis intercept.

The following results were obtained with compounds according to the invention

<u>Compound of</u> <u>Example No.</u>	<u>Ki (nM)</u>		
	<u>Collagenase</u>	<u>Stromelysin-1</u>	<u>Gelatinase -72KD</u>
2	48800	1510	22.2
3	478	29.7	0.273
5	186	12.6	0.035
6	22000	681	8.42
8	215	11.8	0.059

Description of Specific Embodiments

The invention is further illustrated in the following non-limiting Examples.

In the Examples, the following abbreviations are used:

RT	room temperature
DCCl	N,N'-dicyclohexylcarbodiimide
DMF	dimethylformamide
THF	tetrahydrofuran
TFA	trifluoroacetic acid
hplc	high performance liquid chromatography
Et ₂ O	diethylether
CH ₂ Cl ₂	dichloromethane
EtOAc	ethyl acetate
MeOH	methanol
iPr ₂ O	diisopropyl ether

All temperatures are in °C.

Intermediate 1

3[4-Phenoxybutanoyl]-4(S)-benzyl-2-oxazilidinone

Phenoxybutyric acid (1.22g, 6.8mmol, 1.2 equiv) was heated to reflux with thionyl chloride (0.66ml, 1.07g, 9.0mmol, 1.6 equiv) for 30 mins. The excess thionyl chloride was then removed under vacuum, the residue dissolved in THF (10ml), and added to a cold (-78°C) solution of lithiated (S)-4-benzyl-2-oxazilidinone [prepared from the oxazilidinone (1g, 5.6mmol) and n-butyl lithium (1.6M solution, 4.23ml, 6.8mmol, 1.2 equiv) in THF (25ml)]. The reaction mixture was stirred for 2 hrs at -78°C, then quenched with a 1:1 mixture of saturated brine and 10% HCl acid (20ml). The mixture was allowed to warm to room temperature and partitioned between EtOAc and water. The layers were separated and the aqueous layer washed twice with EtOAc. The combined organic layers were washed once with brine, once with sodium bicarbonate solution, dried over MgSO₄ and the solvent removed under vacuum to give a brown oil which was recrystallised from EtOAc to give the title compound as a white solid (0.897g).

¹H NMR CDCl₃ δ: 7.35-7.17 (m, 7H); 6.94-6.86 (m, 3H); 4.66 (m, 1H); 4.18 (m, 2H); 4.06 (t, 2H); 3.30 (dd, 1H); 3.16 (t, 2H); 2.25 (dd, 1H); 2.18 (m, 2H).

Intermediate 2

3-[1-oxo-2(R)-[t-butyl-acetyl-5-(phenoxy)propyl]-4(S)-benzyl-2-oxazolidinone

Sodium hexamethyldisilazide (1.0M solution in THF, 4.2ml, 4.2mmol, 1.4 equiv) was added dropwise to a solution of Intermediate 1. (1.01g, 3mmol) in THF (25ml) at -78°C. Stirring was continued at this temperature for 1hr, then t-butyl bromoacetate (1.74g, 1.44ml, 9mmol, 3 equiv) was added. The temperature was allowed to rise to -20°C over a period of 4 hrs. The reaction was then cooled to -78°C and quenched with a mixture of brine and 10% HCl acid (1:1, 20ml). The reaction was warmed to room temperature and partitioned between water and EtOAc. The aqueous layer was washed twice with EtOAc and the combined organic layers washed once with sodium bicarbonate solution, once with brine, dried over MgSO₄ and the solvent removed in vacuo to give a yellow oil, which was purified on Silica gel (Merck 9385) eluting with 25% EtOAc/hexane to give the title compound as a white solid, (0.90g).

¹H NMR CDCl₃ δ: 7.40-7.18 (m, 7H); 6.95-6.80 (m, 3H); 4.62 (m, 1H); 4.42 (m, 1H); 4.28-3.96 (m, 3H); 3.87 (t, 1H); 3.33 (dd, 1H); 2.91 (dd, 1H); 2.72 (dd, 1H); 2.56 (dd, 1H); 2.26-1.94 (m, 2H); 1.42 (s, 9H).

Intermediate 3

2-(R)-[3-(2-Phenoxyethyl)]succinic acid-4-t-butyl monoester

Hydrogen peroxide (0.91ml, 8mmol, 4 equiv 27.5 wt%) was added to a cold (0°C) solution of Intermediate 2 (0.9g, 2mmol) in THF/water (4:1, 50ml). The solution was stirred for 5 mins then a solution of lithium hydroxide (80mg, 2mmol) in water (10ml) was added dropwise. The reaction was stirred for 2 hours at 0°C then most of the THF was removed in vacuo. The mixture was partitioned between water and CH₂Cl₂, and the aqueous layer separated, acidified to pH1 with 10% HCl acid and extracted with EtOAc (x3). The combined organic layers were dried over MgSO₄ and the solvent removed to give the title compound as a colourless oil, (0.60g, quantitative), which was used without further purification.

Intermediate 4

L- β -cyclohexylalanine-N-(2-phenylethyl) amide

t-Boc- β -cyclohexyl-L-alanine (1.35g, 5mmol) was dissolved in dry CH_2Cl_2 . 4-Nitrophenol (695mg, 5mmol) was added followed by DCCl (1.03g, 5mmol). After 1 hour at room temperature the reaction was concentrated in vacuo, Et_2O was added and the solution filtered. The residue was concentrated in vacuo, dissolved in CH_2Cl_2 (10ml) and phenethylamine (690 μl , 5.5mmol) was added. The reaction was poured into sodium bicarbonate solution and extracted with CH_2Cl_2 (3x20ml), was dried (Na_2SO_4) and concentrated in vacuo.

Purification on silica gel (Merck 9385) using $\text{CH}_2\text{Cl}_2 \rightarrow \text{CH}_2\text{Cl}_2/\text{MeOH}$ (85:15) gave a clean oil (900mg) which was concentrated in vacuo dissolved in CH_2Cl_2 (50ml) and poured into sodium bicarbonate solution. The organic layer was separated, dried (Na_2SO_4) and concentrated in vacuo to give an oil which was purified on silica gel (Merck 9385) using $\text{CH}_2\text{Cl}_2/\text{MeOH}/\text{triethylamine}$ (96:3:1) to give the title compound as an oil (500mg).

^1H NMR CDCl_3 δ : 0.95 (m, 2H); 1.25 (m, 6H); 1.55 (br s, 2H); 1.65 (m, 5H); 2.8 (t, 2H, J=6HZ); 3.4 (dd, 1H, J=3 and 10HZ); 3.5 (dd, 2H, J=6 and 12 HZ); 7.2 (m, 5H).

Intermediate 5

Methyl-4-(4-pyridyl)-buta-1,3-diene carboxylate

Sodium hexamethyldisilazide (23ml, 1.0M solution in THF, 23 mmol, 1.2 equiv) was added to a cold (0°C) solution of trimethyl phosphonocrotonate (4.00g, 19.2mmol) in THF (60ml) under a N_2 atmosphere. The solution was warmed to room temperature during which time a gummy brown precipitate was formed. After stirring for 30 mins the reaction was cooled to 0°C and 4-pyridine carboxaldehyde (2.06g, 1.83ml, 19.2mmol, 1 equiv) was added

dropwise. The reaction was warmed to room temperature and stirred for 4h. The reaction was then poured into ice water and extracted seven times with EtOAc. The combined organic layers were dried over MgSO_4 , the solvent removed under vacuum to give a red/brown gummy solid which was recrystallised from MeOH/water to give the title compound (1.01g).

$^1\text{Hnmr}$, CD_3OD δ : 8.51 (d, 2H, CHN); 7.53 (d, 2H, CHCHN); 7.46-7.22 (m, 2H, CH=); 6.95 (d, 2H); 6.18 (d, 2H); 3.76 (s, 3H).

Intermediate 6

Methyl-4-(4-pyridyl)-butane carboxylate

Palladium on carbon (ca 30mg) was added to a degassed solution of Intermediate 5 (1.94g, 10.3mmol) in MeOH (20ml) and stirred under a H_2 atmosphere for 12h. The reaction mixture was filtered through a silica pad and the residue was washed several times with MeOH. The washings were combined and the solvent removed under vacuum to give the title compound as a brown oil (1.94g).

$^1\text{Hnmr}$ CD_3OD δ : 8.41 (dd, 2H); 7.28 (dd, 2H); 3.64 (s, 3H); 2.67 (t, 2H); 2.37 (t, 2H); 1.68 (m, 4H).

Intermediate 7

5-(4-pyridyl)valeric acid

Lithium hydroxide monohydrate (1.26g, 30.1mmol, 3 equiv) was added to a solution of Intermediate 6 (1.94g, 10.1 mmol) in MeOH (25ml) and stirred at room temperature for 4h under a N_2 atmosphere. After this time the solution was acidified to pH5 with 10% HCl acid and the off white precipitate removed by filtration, washed once with methanol, twice with Et_2O and dried to give the title compound as an off white powder (1.24g).

Intermediate 8**(S)-3-[1-oxo-5-(4-pyridyl)pentyl]-4-(phenylmethyl)-2-oxazolidinone**

A solution of Intermediate 7 (0.63g, 3.52 mmol), (S)-4-phenylmethyl-2-oxazolidinone (0.56g, 3.17 mmol), DCCl (0.91g, 4.4mmol, 1.25 equiv) and N,N-dimethylaminopyridine (0.22g, 1.8 mmol, 0.5 equiv) was heated to reflux for 18h under N₂. After this time the reaction mixture was filtered and the solvent removed under vacuum. The product was purified on silica gel (Merck 9385) eluting with EtOAc/hexane (95:5) to give the title compound as a colourless oil (0.71g) .

¹Hnmr CD₃OD δ: 8.40 (dd, 2H, J=1.5, 4.5Hz); 7.32-7.18 (m, 7H); 4.72 (m, 1H); 4.32-4.18 (m, 2H); 3.11 (dd, 1H, J=3.3, 13.5 Hz); 2.91 (m, 3H); 2.70 (m, 2H); 1.72 (m, 4H).

Intermediate 9**3-[1-oxo-2(R)-(t-butylacetyl)-5-(4-pyridyl)pentyl]-4-(S)-phenylmethyl-2-oxazolidinone**

Sodium hexamethyldisilazide (1.95ml of 1M solution in THF, 1.95mmol, 1.2 equiv) was added dropwise to a solution of Intermediate 8 (0.55g, 1.63mmol) in THF (25ml) at -78°C under N₂. The reaction was stirred at -78°C for 30 mins then a solution of t-Butylbromoacetate (0.32g, 0.26ml, 1.63mmol, 1 equiv) in THF (5ml) was added dropwise and the reaction stirred at -78°C for 4h, then warmed to -20°C for 30 mins. The reaction was quenched at -78°C with a saturated NH₄Cl solution (10ml). The reaction was warmed to room temperature and partitioned between EtOAc and water. The aqueous layer was extracted twice with EtOAc then the combined organic layers washed with brine, dried over MgSO₄ and the solvent removed to give 0.76g of a yellow oil, which was purified in silica gel (Merck 9385) eluting with EtOAc (100%) to give the title compound as a colourless oil (0.53g)

¹Hnmr CD₃OD δ: 8.38 (dd, 2H, J=1.35, 4.6); 7.33-7.21 (m, 7H); 4.69 (m, 1H); 4.26-4.12 (m, 3H); 3.14 (dd, 1H, J=3.2, 13.5Hz); 2.89-2.63 (m, 4H); 2.49 (dd, 1H, J=4.7, 16.5Hz); 1.79-1.57 (m, 4H); 1.42 (s, 9H).

Intermediate 10

2-(R)-[3-(4-pyridyl)propyl]succinic acid-4-t-butyl monoester

Hydrogen peroxide (0.52ml, 27.5 wt%, 4.69mmol, 4 equiv) was added to a solution of Intermediate 9 (0.53g, 1.17mmol) in THF/water (4:1, 25ml) at 0°C under N₂. Stirred for 5 minutes then a 1M solution of Lithium hydroxide (1.17ml, 1.17mmol, 1 equiv) was added dropwise and stirred at 0°C for 1h. Na₂SO₃ solution (5ml of 1M aqueous solution) was added and the reaction stirred for 5 mins. The bulk of the THF was removed under vacuum and 1M NaOH solution (5ml) added. The aqueous layer was extracted with CH₂Cl₂ then acidified to pH6 with 10% HCl acid and extracted six times with CH₂Cl₂. The combined organic layers were dried over MgSO₄ and the solvent removed under vacuum to give the title compound as a white solid (198mg).

Intermediate 11

{4-t-butoxy-2(R)-[3-(4-pyridyl)propyl]succinyl}-L-β-cyclohexylalanine-N-(2-phenylethyl)amide.

A solution of Intermediate 10 (198mg, 0.68mmol), L-β-cyclohexylalanine-N-(2-phenylethyl)amide (186mg, 0.68mmol), N-methylmorpholine (68.7mg, 74.7μl, 0.68mmol), 1-hydroxybenzotriazole (1.36ml of a 0.5M solution in DMF, 0.68mmol) and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (130.4mg, 0.68mmol) in DMF (10ml) was stirred under N₂ at room temperature for 18h. After this time the reaction was partitioned between EtOAc and phosphate pH7 buffer. The aqueous layer was extracted twice with EtOAc and the combined organic layers washed with brine twice, dried over MgSO₄ and the solvent removed under vacuum to give a yellow oil which was purified on silica gel (Merck 9385) eluting with 5% MeOH/CH₂Cl₂ to give the title compound as a yellowish oil (397mg).

¹Hnmr CHCl₃ δ: 8.42 (dd, 2H); 7.31-6.90 (m, 9H); 4.43 (dd, 1H); 3.58-3.41 (m, 2H); 2.82-2.53 (m, 5H); 1.79-0.75 (m, 19H); 1.42 (s, 9H).

Intermediate 12

{4-t-butoxy-2(R)-[3-(4-pyridyl)propyl]succinyl}-L-β-cyclohexylalanine-N-[2-(p-sulphonamidophenyl)ethyl]amide

By proceeding in a manner similar to Intermediate 11, but replacing the L-β-cyclohexylalanine-N-(2-phenylethyl)amide by L-β-cyclohexylalanine-N-[2-(p-sulphonamidophenyl)ethyl]amide, the title compound was obtained as an off-white powder.

¹Hnmr CD₃OD δ: 7.98 (s, 2H); 7.82 (d, 2H); 7.37 (d, 2H); 7.28 (d, 2H); 4.33 (m, 1H); 3.38 (m, 2H); 3.06-2.54 (m, 6H); 2.32 (dd, 1H); 1.83-0.78 (m 19H); 1.42 (s, 9H).

Example 1

{4-t-Butoxy-2(R)-[3-(2-phenoxyethyl)succinyl]-L-β-cyclohexylalanine-N-(2-phenylethyl)amide

Ethyl chloroformate (196 μl, 222mg, 2.04mmol) 1.05 equiv) was added to a solution of Intermediate 3 (600mg, 2mmol) and N-methylmorpholine (235μl, 216mg, 2.14mmol, 1.1 equiv) in THF (10ml) at -20°C under a N₂ atmosphere. After 1 hour a solution of Intermediate 4 (532mg, 1.94mmol) in THF (5ml) was added and the reaction allowed to warm to room temperature overnight. The reaction mixture was partitioned between EtOAc and water. The aqueous layer was extracted twice with EtOAc and then the combined organic layers were washed once with 10% HCl acid, once with brine and once with saturated sodium bicarbonate solution, dried over MgSO₄ and the solvent removed to give an oil which was purified on silica gel (Merck 9385) eluting with 1% MeOH/CH₂Cl₂ to give the title compound as a colourless oil (814mg).

¹H NMR CDCl₃ δ: 7.33-6.87 (m, 10H); 6.18 (m, 2H); 4.35 (m, 1H); 3.96 (m, 2H); 3.41 (m, 2H); 2.92-2.66 (m, 3H); 2.43 (dd, 1H); 1.80-0.78 (m, 15H); 0.96 (s, 9H).

Example 2

[4-Hydroxy-2-(R)-[3-(2-phenoxyethyl)succinyl]-L-β-cyclohexylalanine-N-(2-phenylethyl)amide

The compound of Example 1 (0.80g, 1.45mmol) was heated with water (1ml) and TFA (4ml) and stood overnight (18hrs) at room temperature. The volatiles were removed under vacuum and the product was purified on silica gel (Merck 9385) eluting with EtOAc/hexane (1:1) to give 0.50g of a powdery solid. 100mgs of this were further purified by reverse phase hplc using TFA/H₂O/MeCN (starting with 0.1:50:50 ending with 0.1:0:100 over 20 minutes) to give 30mgs of the title compound as a white solid.

¹H NMR CD₃OD δ: 7.26-7.13 (m, 7H); 6.92-6.87 (m, 3H); 4.36 (dd, 1H); 3.97-3.93 (m, 2H); 2.99 (m, 1H); 2.72-2.65 (m, 3H); 2.50 (dd, 1H, J=16.7, 5.0Hz); 2.03-0.83 (m, 17H).

Example 3

4-(N-Hydroxyamino)-2-(R)-[3-(2-phenoxyethyl)succinyl]-L-β-cyclohexylalanine-N-(2-phenylethyl)amide

Ethyl chloroformate (122g, 108μl), 1.1mmol, 1.4 equiv) was added to a -20°C solution of the compound of Example 2 (398mg 0.8mmol) and N-methylmorpholine (123mg, 132μl, 1.2mmol, 1.5 equiv) in THF (10ml). The white suspension was stirred at -20°C for 1 hr then O-trimethylsilylhydroxylamine (169mg, 197μl, 1.6mmol, 2 equiv) was added and the solution allowed to warm to room temperature overnight. The reaction was partitioned between EtOAc and 10% HCl acid. The aqueous layer was washed twice with EtOAc and the combined organic layers washed once with brine and once with sodium bicarbonate solution, dried over MgSO₄ and the solvent removed under vacuum to give a white solid (316mg), which was purified by reverse phase hplc using TFA/water/MeCN (starting with 0.1:50:50 and ending with 0.1:0:100 over 20 mins) to give the title compound as a white solid (142mg)

^1H NMR CD_3OD δ : 7.25-7.13 (m, 7H); 6.91-6.87 (m, 3H); 4.32 (dd, 1H); 3.97-3.91 (m, 2H); 3.31 (m, 1H); 2.70 (t (app) $J=7.2\text{Hz}$, 2H); 2.42 (dd, 1H); 2.27 (dd, 1H); 2.04-0.87 (m, 17H).

Example 4

{4-Hydroxy-2(R)-[3-(4-pyridinium)propyl]succinyl}-L- β -cyclohexylalanine-N-(2-phenylethyl)amide trifluoroacetate

A solution of Intermediate 11 (397mg, 0.60ml) in TFA (4ml) and water (1ml) was stood at room temperature for 3h. The volatiles were then removed under vacuum to give an oil which was purified on silica gel (Merck 9385) eluting with $\text{CH}_2\text{Cl}_2/\text{MeOH}$ (9:1) to give the title compound as a colourless oil (328g).

About 100mg were further purified by reverse phase hplc eluting with $\text{TFA}/\text{H}_2\text{O}/\text{MeOH}$ (starting with 0.1:60:40) and ending with 0.1:40:60 over 20 mins).

^1H nmr CD_3OD δ : 8.65 (d, 2H, $J=6.4\text{Hz}$); 7.91 (d, 2H, $J=6.5\text{Hz}$); 7.28-7.15 (m, 5H); 4.36 (dd, 1H, $J=5.3, 9.6\text{Hz}$); 3.49-3.28 (m, 2H); 2.99-2.59 (m, 4H); 2.39 (d, 1H, $J=5.2, 16.9\text{Hz}$); 1.84-0.74 (m, 19H).

^{13}C nmr CD_3OD δ : 176.8, 175.1, 174.9, 142.3, 140.4, 129.8, 129.5, 128.4, 127.4, 52.5, 41.9, 40.6, 37.6, 36.5, 36.4, 35.0, 34.8, 33.3, 33.1, 28.1, 27.6, 27.3, 27.0.

Example 5

{4-(N-hydroxyamino)-2(R)-[3-(4-pyridinium)propyl]succinyl}-L- β -cyclohexylalanine-N-(2-phenylethyl)amide trifluoroacetate

Ethylchloroformate (84.8mg, 74.7 μl , 0.78mmol, 2.1 equiv) was added to a cold (-20°C) solution of the compound of Example 4 (225mg, 0.37mmol) and N-methyl morpholine (93.6mg, 101.7 μl , 0.93mmol, 2.5 equiv) in THF

(5ml) under N₂. Stirring was continued at -20°C for 1h then O-trimethylsilyloxyhydroxylamine (172mg, 200μl, 17mmol, 4.6 equiv) was added and the reaction allowed to warm to room temperature and stirred overnight to give a white suspension, which was poured into saturated NH₄Cl solution and extracted 3 times with EtOAc. The combined organic layers were washed once with brine, dried over MgSO₄ and solvent removed under vacuum to give a yellowish solid which was triturated with CH₂Cl₂/iPr₂O to give the title compound as an off white solid (225gm).

About 70mg were further purified by reverse phase hplc eluting with TFA/H₂O/MeCN (starting at 0.1:70:30 and ending with 0.1:50:50 over 20 mins) to give the title compound as a hygroscopic white solid (26.5mg).

¹Hnmr δ CD₃OD: 8.66 (br s, 2H); 7.88 (d, 2H); 7.30-7.15 (m, 5H); 4.32 (dd, 1H); 3.49-3.34 (m, 2H); 3.03-2.71 (m, 5H); 2.41-2.16 (m, 2H); 1.82-0.81 (m, 19H).

Example 6

{4-(N-hydroxyamino)-2-(R)-[3-(N-methyl-4-pyridinium)propyl]succinyl}-L-β-cyclohexylalanine-N-(2-phenylethyl)amide iodide

The compound of Example 5 (48.0mg, 0.08mmol) was dissolved in 1ml methyl iodide and stirred at room temperature for 18h. The excess methyl iodide was removed by evaporation with N₂ and the resulting yellow gum was purified by reverse phase hplc eluting with TFA/H₂O/CH₃CN (starting with 0.1:70:30 and ending with 0.1:50:50 over 20 mins), to give the title compound as a hygroscopic gum (25mg).

¹Hnmr CD₃OD δ: 8.61 (d, 2H); 7.84 (d, 2H); 7.25-7.10 (m, 2H); 4.26 (m, 1H); 4.21 (s, 3H); 3.44-3.22 (m, 2H); 2.96-2.65 (m, 4H); 2.34-2.07 (m, 2H); 1.80-0.75 (m, 19H).

Example 7**[4-Hydroxy-2(R)-[3-(4-pyridinium)propyl]succinyl]-L-β-cyclohexylalanine-N-[2-(p-sulphonamidophenyl)ethyl]amide trifluoroacetate**

By proceeding in a manner similar to Example 4 with Intermediate 12, the title compound was obtained as an off-white hygroscopic foam.

¹Hnmr CD₃OD δ: 8.68 (d, 2H); 7.95 (d, 2H); 7.80 (d, 2H); 7.48 (d, 2H); 4.36 (m, 1H); 3.46 (m, 2H); 3.18-2.80 (m, 5H); 2.62 (dd, 1H); 2.51 (dd, 1H); 1.87-0.80 (m, 19H).


Example 8**{4-(N-Hydroxyamino)-2-(R)-[3-(4-pyridinium)propyl]succinyl]-L-β-cyclohexylalanine-N-[2-(p-sulphonamidophenyl)ethyl]amide trifluoroacetate**

A solution of the compound of Example 7 (0.30g, 0.44mmol) in THF (5ml) was treated with N-methylmorpholine (111mg, 120μl, 1.10mmol), and cooled to -20°C. Ethyl chloroformate (50.4mg, 44.4μl, 0.46mmol) was added and the resulting slurry stirred for 15 mins at -20°C. O-trimethylsilyl hydroxylamine (230mg, 267μl, 2.2mmol) was then added and the reaction allowed to warm to room temperature over 18 hrs. The mixture was partitioned between EtOAc and phosphate pH7 buffer. The aqueous layer was extracted with EtOAc (x5) and the combined organic layers washed with brine then dried over MgSO₄, and solvent removed under vacuum to give a whitish solid, which was purified by reverse phase hplc (gradient elution over 20 mins TFA:water:MeCN 0.1:70:30 to 0.1:60:40) to give the title compound as a hygroscopic white solid (46 mg).

¹Hnmr CH₃OD δ: 8.52 (br s, 2H); 8.14 (br d, 1H); 7.83 (br s, 1H); 7.75 (d, 2H); 7.47 (d, 2H); 4.31 (m, 1H); 3.46 (m, 2H); 3.04-2.74 (m, 6H); 2.38 (d, 1H); 2.20 (d, 1H); 8.2-0.80 (m, 19H).

CLAIMS

1. A compound of formula (1)



wherein R represents a -CONHOH, carboxyl (-CO₂H), esterified carboxyl or -P(O)(X¹R⁸)X²R⁹ groups, where X¹ and X², which may be the same or different is each an oxygen or a sulphur atom, and R⁸ and R⁹, which may be the same or different each represents a hydrogen atom or an optionally substituted alkyl, aryl or aralkyl group;

R¹ represents a hydrogen atom or an optionally substituted alkyl, alkenyl, aryl, aralkyl, heteroaralkyl or heteroarylthioalkyl group;

R² represents an optionally substituted aryloxy, arylthio, aryloxyalkyl, arylthioalkyl, heteroaryl, heteroaralkyl, heteroaryloxy, heteroarylthio, heteroaryloxyalkyl or heteroarylthioalkyl group;

R³ represents a hydrogen atom or an alkyl group;


R⁴ represents a hydrogen atom or an alkyl group;

R⁵ represents an optionally substituted alkyl or alkenyl group optionally interrupted by one or more -O- or -S- atoms or -N(R⁷) groups, [where R⁷ is a hydrogen atom or a C₁₋₆ alkyl group] or a group -[Alk]_nR⁶ where Alk is an alkyl or alkenyl group optionally interrupted by one or more -O- or -S- atoms or -N(R)⁷ groups, n is zero or an integer 1, and R⁶ is an optionally substituted cycloalkyl or cycloalkenyl group;

X represents an amino (-NH₂), or substituted amino, hydroxyl or substituted hydroxyl group;

and the salts solvates and hydrates thereof.

2. A compound according to Claim 1 wherein R represents a-CONHOH group.
3. A compound according to Claim 1 or 2 wherein R¹, R³ and R⁴ is each a hydrogen atom.
4. A compound according to any of Claims 1 to 3 wherein R² is an aryloxyalkyl, heteroaralkyl or heteroaryloxyalkyl group.
5. A compound according to Claim 4 wherein R² is an optionally substituted phenoxyalkyl, pyridinylalkyl or pyridinyloxyalkyl group.
6. A compound according to Claims 1 to 5 wherein R⁵ is a C₁₋₆ alkyl group or an AlkR⁶ group, where Alk is a C₁₋₆ alkyl group and R⁶ is a cycloalkyl or cycloalkenyl group.
7. A compound according to any of the preceding claims where X is an amino or substituted amino group.
8. A compound of formula (1a)



wherein R represents a CONHOH, carboxyl (-CO₂H) esterified carboxyl or -P(O) (X¹R⁸)X²R⁹ group, where X¹ and X², which may be the same or different is each an oxygen or a sulphur atom, and R⁸ and R⁹, which may be the same or different each represents a hydrogen atom or an optionally substituted alkyl, aryl or aralkyl group;

R² represents an optionally substituted aryloxy, arylthio, aryloxyalkyl,

arylthioalkyl, heteroaryl, heteroaralkyl, heteroaryloxy, heteroarylthio, heteraryloxyalkyl or heterarylthioalkyl group;

R⁵ represents an optionally substituted alkyl or alkenyl group optionally interrupted by one or more -O- or -S- atoms or -N(R⁷) groups, [where R⁷ is a hydrogen atom or a C₁₋₆ alkyl group] or a group -[Alk]_nR⁶ where Alk is an alkyl or alkenyl group optionally interrupted by one or more -O- or -S- atoms or -N(R)⁷ group, n is zero or an integer 1, and R⁶ is an optionally substituted cycloalkyl or cycloalkenyl group;

X represents an amino (-NH₂), or substituted amino, hydroxyl or substituted hydroxyl group; and the salts, solvates and hydrates thereof;

9. 4-t-Butoxy-2(R)-[3-(2-phenoxyethyl)succinyl]-L-β-cyclohexylalanine-N-(2-phenylethyl)amide;

4-Hydroxy-2(R)-[3-(2-phenoxyethyl)succinyl]-L-β-cyclohexylalanine-N-(2-phenylethyl)amide;

4-(N-Hydroxyamino-2(R)-[3-(2-phenoxyethyl)succinyl]-L-β-cyclohexylalanine-N-(2-phenylethyl)amide;

{4-Hydroxy-2(R)-[3-(4-pyridinium)propyl)succinyl]-L-β-cyclohexylalanine-N-(2-phenylethyl)amide;


{4-(N-Hydroxyamino)-2(R)-[3-(4-pyridinium)propyl)succinyl]-L-β-cyclohexylalanine-N-(2-phenylethyl)amide;

{4-(N-Hydroxyamino-2(R)-[3-(N-methyl-4-pyridinium)propyl)succinyl]-L-β-cyclohexylalanine-N-(2-phenylethyl)amide; and the salts, solvates and hydrates thereof.


10. A pharmaceutical composition comprising a compound according to any one of Claims 1 to 9 and a pharmaceutically acceptable diluent, carrier or excipient.

11. A process for preparing a compound of formula (1) as defined in Claim 1, the process comprising:

- (a) coupling an acid of formula (2)




or an active and/or protected derivative thereof, with an amine of formula (3)



followed by removal of any protecting groups.

- (b) oxidising and oxazolidinone of formula (5)



(where Ph is a phenyl group)

- (c) interconverting a compound of formula (1) to another compound of formula (1).

INTERNATIONAL SEARCH REPORT

PCT/GB 93/01186

International Application No

I. CLASSIFICATION OF SUBJECT MATTER (if several classification symbols apply, indicate all) ⁶		
According to International Patent Classification (IPC) or to both National Classification and IPC		
Int.Cl. 5 C07C237/22; C07C259/06; C07D213/56; A61K31/16 A61K31/44		
II. FIELDS SEARCHED		
Minimum Documentation Searched ⁷		
Classification System	Classification Symbols	
Int.Cl. 5	C07C ; C07D	
Documentation Searched other than Minimum Documentation to the extent that such Documents are included in the Fields Searched ⁸		
III. DOCUMENTS CONSIDERED TO BE RELEVANT⁹		
Category ¹⁰	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³
A	EP,A,0 214 639 (SEARLE, G. D., AND CO.) 18 March 1987 cited in the application see abstract; claims; examples 13-15,17 ---	1-11
A	EP,A,0 274 453 (LABORATOIRE ROGER BELLON) 13 July 1988 cited in the application see claims; examples ---	1-11
A	WO,A,9 005 716 (BRITISH BIO-TECHNOLOGY) 31 May 1990 cited in the application see page 1, line 7 - line 29; claims; examples ---	1-11
A	WO,A,9 115 507 (BEECHAM) 17 October 1991 see claims ---	1-11
		-/--
<p>⁹ Special categories of cited documents : ¹⁰</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"Δ" document member of the same patent family</p>		
IV. CERTIFICATION		
Date of the Actual Completion of the International Search 13 SEPTEMBER 1993		Date of Mailing of this International Search Report 21. 09. 93
International Searching Authority EUROPEAN PATENT OFFICE		Signature of Authorized Officer SEUFERT G.H.

III. DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)		
Category *	Citation of Document, with indication, where appropriate, of the relevant passages	Relevant to Claim No.
P,A	EP,A,0 489 577 (CELLTECH) 10 June 1992 see page 2, line 29 - line 31; claims; examples -----	1-11

INTERNATIONAL SEARCH REPORT

International application No.

PCT/GB93/01186

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
The vast number of theoretically conceivable compounds resulting from the combination of all claimed substituents precludes a comprehensive search for compound (1). For economic reasons the search has been limited. Guided by the spirit of the application and the inventive concept disclosed in the ./.

3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.

2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.

3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:

4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/

in the descriptive part the search has been limited to the following cases:

$R^1, R^3, R^4 = H$;

$R^5 =$ substituted alkyl

$R = CONHOH, CO_2H, CO_2R$

$X =$ subst. amino

$R^2 =$ as defined in the appl. with alkyl = C_{0-6}

(Guidelines Exam. Part B, Chap. III 3.6, 3.7)

**ANNEX TO THE INTERNATIONAL SEARCH REPORT
ON INTERNATIONAL PATENT APPLICATION NO.**

GB 9301186
SA 75173

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report.
The members are as contained in the European Patent Office EDP file on

The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information. 13/09/93

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A-0214639	18-03-87	US-A- 4599361	08-07-86
		US-A- 4743587	10-05-88
		AU-B- 588362	14-09-89
		AU-A- 6240886	12-03-87
		JP-A- 62103052	13-05-87
EP-A-0274453	13-07-88	FR-A- 2609289	08-07-88
		JP-A- 63258449	25-10-88
		US-A- 4918105	17-04-90
WO-A-9005716	31-05-90	AU-A- 4746890	12-06-90
		EP-A- 0445206	11-09-91
		JP-T- 4503057	04-06-92
WO-A-9115507	17-10-91	AU-B- 634383	18-02-93
		AU-A- 7671191	30-10-91
		EP-A- 0527764	24-02-93
EP-A-0489577	10-06-92	AU-A- 9017391	25-06-92
		AU-A- 9023391	25-06-92
		EP-A- 0489579	10-06-92
		WO-A- 9209564	11-06-92
		WO-A- 9209565	11-06-92
		GB-A- 2255339	04-11-92
		GB-A- 2255340	04-11-92
		JP-T- 5503719	17-06-93
		JP-T- 5503720	17-06-93